

Morphic cohomology of toric varieties

Abdó Roig-Maranges*

January 18, 2010

In this paper we construct a spectral sequence computing a modified version of morphic cohomology of a toric variety (even when it is singular) in terms of combinatorial data coming from the fan of the toric variety.

Contents

1	Introduction	2
2	Morphic cohomology	2
2.1	Bivariant Morphic spaces	4
2.2	Morphic cohomology complexes	7
2.3	A modification of the morphic cohomology	9
2.4	Some remarks about notation	12
2.5	Cup product and Kunneth homomorphism	13
3	Toric varieties	15
3.1	Cones and fans	16
3.2	Definition of a toric variety	17
3.3	Morphic cohomology of an algebraic torus	20
4	Spectral sequence associated to a toric variety	21
4.1	Resolution associated to a fan	21
4.2	The spectral sequence	26
4.3	Some applications	27

*Partially supported by MICINN MTM2009-09557 from the Spanish government and the “Comissionat per a Universitats i Recerca” from the “Generalitat de Catalunya”.

1 Introduction

Morphic cohomology is a cohomological theory on algebraic varieties introduced by Friedlander and Lawson in [FL92]. On one hand this theory has a very geometric definition, and on the other it is strongly related to the abstractly defined motivic cohomology theory.

In this paper, we define, along the lines of [Fri00], a modification of morphic cohomology in order to have Mayer-Vietoris and homotopy invariance properties of the theory for all quasi-projective varieties (not only smooth). Then, we use those properties to construct a spectral sequence computing the morphic cohomology of a toric variety in terms of its combinatorial data.

In section 2 we recall the definition of morphic cohomology and the theorems we need, and define the modification of the theory we will use. This is mainly an expository section except, maybe, subsection 2.3.

In section 3 we recall the definition of toric variety, and the constructions we will use, mainly to set up the notation. We also write down an explicit computation of the morphic cohomology of algebraic tori.

Finally, in section 4 we build a resolution of the constant sheaf $\mathbb{Z}_X$ on a toric variety in terms of the combinatorial data. This resolution allows us to construct a spectral sequence (theorem 4.21) computing the hypercohomology of a complex of sheaves $\mathcal{F}^*$ on the toric variety $X(\Delta)$ in terms of the combinatorial data and the value of this hypercohomology on algebraic tori. Then we specialize this to the case of morphic cohomology, giving a very explicit spectral sequence converging to the morphic cohomology of $X(\Delta)$ and whose second page involves only combinatorial data. Moreover, we prove its rational degeneration (theorem 4.23). Finally, we also use the spectral sequence to extend the Suslin conjecture, proved in [FHW04] and [Voi08b] for smooth linear varieties, to all toric varieties, even singular ones.

I want to express my gratitude to my advisor, Pere Pascual, for his help and encouragement during this work, and also to Eric Friedlander and Mircea Voineagu, from whom I learnt a lot during the time I spent in Los Angeles.

2 Morphic cohomology

In this section we recall some facts about morphic cohomology. Let Y be a projective variety over $\mathbb{C}$, with a fixed projective embedding. It is a classical fact (see chapter 1 in [Kol96]) that the set of effective k -cycles on Y has the structure of an algebraic variety. This variety is called the *Chow variety*, and we will denote it by $C(Y, k)$. It may have infinitely many connected components, corresponding to the homology classes of the cycles. The Chow variety has a natural operation given by the sum of cycles. This operation is algebraic, so $C(Y, k)$ is a monoid in the category $\mathbf{Proj}_{\mathbb{C}}$ of projective varieties (see [Fri91]).

Set theoretically, the group of algebraic k -cycles on Y is the group completion of the Chow variety,

$$Z(Y, k) = C(Y, k)^+ = C(Y, k) \times C(Y, k) / C(Y, k). \quad (1)$$

This can be used to induce a topology on $Z(Y, k)$ from the analytic topology on the Chow variety. As shown in [Law89] and [Fri91], this topology on $Z(Y, k)$ does not depend on the projective embedding of Y . Moreover, its homotopy type is a very interesting invariant containing information about the geometry of Y . The homotopy groups of this space $Z(Y, k)$ are called the *Lawson homology groups* of Y , and are denoted by $L_k H_n(Y)$. The usual indexing convention is as follows,

$$L_k H_n(Y) = \pi_{n-2k} Z(Y, k). \quad (2)$$

The construction of the Chow varieties $C(Y, k)$ works only for projective Y , but the topology on $Z(Y, k)$ can be defined for quasi-projective varieties (see [LF92]). Take $Y \subset \bar{Y}$ a projective closure of the quasi-projective variety Y , and let $Y^\infty = \bar{Y} \setminus Y$, which is again projective. Then we define the topology in $Z(Y, k)$ as the quotient topology given by the set-theoretic identification

$$Z(Y, k) = Z(\bar{Y}, k) / Z(Y^\infty, k). \quad (3)$$

Theorem 4.3 in [LF92] states that the topology on $Z(Y, k)$ we have just described does not depend on the choice of projective closure $\bar{Y}$.

It will be useful to work simplicially, so we denote the singular simplicial set associated to $Z(Y, k)$ by

$$Z_\bullet(Y, k) = \text{Sing}_\bullet Z(Y, k).$$

The group law on $Z(Y, k)$ makes $Z_\bullet(Y, k)$ into a simplicial abelian group.

One important fact about those spaces of cycles is that the homotopy groups of the spaces of 0-cycles coincide with the Borel-Moore homology groups of Y (singular homology if Y is projective),

$$\pi_l Z(Y, 0) \simeq H_l^{\text{BM}}(Y, \mathbb{Z}). \quad (4)$$

This is a consequence of a classical result of Dold and Thom [DT58]. From this we see that the space of cycles $Z(\mathbb{A}^q, 0)$ has only nontrivial homotopy groups in dimension $2q$, so it is an algebraic model of an Eilenberg-MacLane space $K(\mathbb{Z}, 2q)$.

This interpretation motivates the construction of a cohomological theory related to the Lawson homology groups, *morphic cohomology*, in the same way singular cohomology is related to singular homology. The inspiration for constructing morphic cohomology comes from the fact that singular cohomology can be interpreted as homotopy groups of spaces of continuous maps to an Eilenberg-MacLane space,

$$H^{2q-l}(X, \mathbb{Z}) \simeq \pi_l \text{Map}(X, K(\mathbb{Z}, 2q)). \quad (5)$$

Then one could think of the homotopy groups of spaces of “algebraic maps” from an algebraic variety X to our algebraic model of $K(\mathbb{Z}, 2q)$, the space of 0-cycles $Z(\mathbb{A}^q, 0)$. This is roughly

the idea behind the construction of morphic cohomology, as we will recall in section 2.2. But before this, it will be convenient to talk about a more general bivariant theory depending on two algebraic varieties, X and Y .

2.1 Bivariant Morphic spaces

We are going to describe a bivariant theory, following [Fri98]. This theory depends on two algebraic varieties X and Y , and roughly speaking, is defined as the homotopy groups of spaces of “algebraic maps” from X to a space of cycles $Z(Y, k)$. The first thing to do is to make precise what we mean by “algebraic maps”. For technical reasons, it happens to be more convenient to allow a slightly more general class of maps than the usual algebraic morphisms of varieties.

Definition 2.1. A *continuous algebraic map* between two algebraic varieties X and Y is a rational map $f: X \dashrightarrow Y$ which is defined at every point and is continuous. We will denote by $\text{Mor}(X, Y)$ the set of continuous algebraic maps.

Remark 2.2. If X is a normal variety, a continuous algebraic map $f: X \rightarrow Y$ is a morphism. This is a consequence of Zariski’s main theorem applied to the projection $\Gamma_f \rightarrow X$ of the graph of the map to X .

In particular, this means that the technical point involving continuous algebraic maps appears only for non-normal varieties.

For more details on the role of continuous algebraic maps see [Fri00] section 1 and [FW01].

We want to topologize the spaces $\text{Mor}(X, Y)$. The first choice of topology would be the compact-open topology, however it is not the right one. Its problems come from two different sources: one appears when X is non-complete, and the other when X is non-normal. Here we just describe the correct topology we need, and for the justification of why this seems to be the right choice we refer to [FL97] appendix C (for the case in which X is normal) and [Fri00] proposition 1.4 for the general case.

Definition 2.3. Let X be a quasi-projective variety and Y a projective one.

When X is normal, we provide the space $\text{Mor}(X, Y)$ of continuous algebraic maps with a topology such that a sequence of maps $\{f_n\}$ converges to $f \in \text{Mor}(X, Y)$ if, and only if

1. it converges for the compact-open topology,
2. there is a compactification $X \subset \overline{X}$ for which the sequence of closures of graphs $\overline{\Gamma_{f_n}} \subset \overline{X} \times Y$ has bounded degree.

When X is not normal, we provide the space $\text{Mor}(X, Y)$ with the topology induced as a subspace of $\text{Mor}(\tilde{X}, Y)$, where $\tilde{X} \rightarrow X$ is the normalization of X .

Remark 2.4. In the case in which X is projective and normal, the condition on the boundeness of the degree becomes void, and then the topology on $\text{Mor}(X, Y)$ coincides with the compact-open topology.

There is one last technical point we must deal with. In the situation in which we are interested, $\text{Mor}(X, Z(Y, k))$, the second argument is the space of cycles $Z(Y, k)$ which is not an algebraic variety. But it is the group completion of the Chow variety $C(Y, k)$, so we can consider instead the well defined space $\text{Mor}(X, C(Y, k))^+$ where $(-)^+$ denotes a group completion as in (1).

Moreover, if Y is quasi-projective, we can perform a similar trick as for $Z(Y, k)$, but on the whole $\text{Mor}(X, C(Y, k))^+$. We chose a projective embedding for Y , and take the projective closure $\bar{Y}$, with closed complement Y^∞ . In this case though, the bivariant theory will depend on the pair $(\bar{Y}, Y^\infty)$, not only on Y . Finally, as it will be convenient to work simplicially, we take singular chains. This is the content of the following definition.

Definition 2.5. Let X be quasi-projective variety and Y a projective one. Then, the *bivariant morphic space* $M_\bullet(X, Y, k)$ is the simplicial abelian group

$$M_\bullet(X, Y, k) = (\text{Sing}_\bullet \text{Mor}(X, C(Y, k)))^+, \quad (6)$$

where the group completion $(-)^+$ here means a levelwise group-completion of the simplicial abelian monoid.

If Y is quasi-projective, with projective closure $\bar{Y}$ and closed complement Y^∞ , Then the *bivariant morphic space* is the quotient simplicial abelian group

$$M_\bullet(X, \bar{Y}/Y^\infty, k) = M_\bullet(X, \bar{Y}, k)/M_\bullet(X, Y^\infty, k). \quad (7)$$

Remark 2.6. The construction of this morphic space is contravariantly functorial in X , by composition. It is also covariantly functorial for proper morphisms of the pair $(\bar{Y}, Y^\infty)$. That is, if $\bar{Y} \rightarrow \bar{Y}'$ is a proper morphism sending Y^∞ to Y'^∞ , the induced map $C(\bar{Y}, k) \rightarrow C(\bar{Y}', k)$ sends the cycles in $C(Y^\infty, k)$ to $C(Y'^\infty, k)$ so we get a continuous map

$$M_\bullet(X, \bar{Y}/Y^\infty, k) \longrightarrow M_\bullet(X, \bar{Y}'/Y'^\infty, k).$$

Those spaces have a localization property with respect to the second argument analogous to proposition 3.2 in [LF92].

Theorem 2.7. Let X be a quasi-projective variety and let $Y_0 \subset Y_1 \subset Y_2$ a triple of projective varieties. Then, the sequence of simplicial abelian groups

$$M_\bullet(X, Y_1/Y_0, k) \rightarrow M_\bullet(X, Y_2/Y_0, k) \rightarrow M_\bullet(X, Y_2/Y_1, k) \quad (8)$$

is a fibration sequence and, in particular, there is an associated long exact sequence of homotopy groups.

Proof. The proof is analogous as proposition 3.2 in [LF92] using the fact that a quotient of simplicial abelian groups produces a fibration sequence (see [GJ99] corollary V.2.7). ●

In the special case in which $X = \text{Spec } \mathbb{C}$ there is the following comparison result

Proposition 2.8. *Let Y be a quasi-projective variety with projective closure $\overline{Y}$ and closed complement Y^∞ . Then the natural map $M_\bullet(\text{Spec } \mathbb{C}, \overline{Y}/Y^\infty) \rightarrow Z_\bullet(Y, k)$ is a weak homotopy equivalence.*

Proof. This follows from [FG93] proposition 1.3 and theorem 1.4, which enable us to commute, up to homotopy equivalence, the singular chains functor with the group completion and the quotient in the definitions of $M_\bullet(X, \overline{Y}/Y^\infty, k)$ and $Z_\bullet(Y, k)$. ●

Remark 2.9. In particular, for a quasi-projective Y with closed subvariety Y_0 and open complement U , theorem 2.7 and proposition 2.8 together with the long exact sequence of homotopy groups associated to a fibration gives the long exact sequence

$$\cdots \longrightarrow L_k H_n(Y_0) \longrightarrow L_k H_n(Y) \longrightarrow L_k H_n(U) \longrightarrow L_k H_{n-1}(Y_0) \longrightarrow \cdots$$

This is the localization theorem for Lawson homology (proposition 4.8 in [LF92]).

Definition 2.10. Let Y be a projective variety with a given projective embedding $Y \subset \mathbb{P}^N$. Let $\mathbb{P}^N \subset \mathbb{P}^{N+1}$ be the embedding in the first N coordinates, and $p^\infty = [0, \dots, 0, 1] \in \mathbb{P}^{N+1}$ (p^∞ is a choice of a point not in $\mathbb{P}^N$). The *Lawson suspension* of Y , denoted by $\mathbb{Z}Y$ is the projective variety in $\mathbb{P}^{N+1}$ of the points contained in a line through p^∞ that meets Y .

Remark 2.11. The Lawson suspension induces a continuous algebraic map at the level of Chow varieties

$$\mathbb{Z}: C(Y, k) \rightarrow C(\mathbb{Z}Y, k+1),$$

such that

$$\mathbb{Z}(n_1[W_1] + \cdots + n_r[W_r]) = n_1[\mathbb{Z}W_1] + \cdots + n_r[\mathbb{Z}W_r].$$

This leads to a very important theorem in the theory.

Theorem 2.12 (Lawson suspension). *The Lawson suspension of cycles induces a map of morphic spaces*

$$M_\bullet(X, \overline{Y}/Y^\infty, k) \longrightarrow M_\bullet(X, \mathbb{Z}\overline{Y}/\mathbb{Z}Y^\infty, k+1) \quad (9)$$

which is a homotopy equivalence.

Proof. See [FL92] theorem 3.3 for the projective case and [Fri98] proposition 3.7 for the quasi-projective version. ●

Corollary 2.13. *The projection $p: X \times \mathbb{A}^1 \rightarrow X$ induces, through pull-back of cycles, a homotopy equivalence*

$$Z_{\bullet}(X, k) \xrightarrow{\sim} Z_{\bullet}(X \times \mathbb{A}^1, k + 1). \quad (10)$$

Proof. Using that $\mathbb{A}^1 \setminus \{p_{\infty}\} = X \times \mathbb{A}^1$, this becomes a consequence of the localization theorem 2.7, the suspension theorem 2.12 and the homotopy equivalence 2.8 between the space of cycles $Z_{\bullet}(Y)$ and $M_{\bullet}(\text{Spec } \mathbb{C}, \overline{Y}/Y^{\infty}, k)$. $\bullet$

2.2 Morphic cohomology complexes

As said in the beginning of this section, to construct the morphic cohomology we will need an algebraic analogue of $\text{Map}(X, K(\mathbb{Z}, 2q))$. Since $Z(\mathbb{A}^q, 0)$ is an algebraic model of a $K(\mathbb{Z}, 2q)$, it would make sense to use the spaces $M_{\bullet}(X, \mathbb{A}^t, k)$. We have defined those spaces for a pair of projective varieties $(\overline{Y}, Y^{\infty})$ not for a quasi-projective Y alone, but $\mathbb{A}^t$ has a natural choice of projective closure, $\overline{Y} = \mathbb{P}^t$ for which the closed complement is

$$Y^{\infty} = \{[x_0 : \cdots : x_t] \in \mathbb{P}^t \mid x_0 = 0\} = \mathbb{P}^{t-1}.$$

So a good candidate spaces computing the morphic cohomology are $M_{\bullet}(X, \mathbb{P}^t/\mathbb{P}^{t-1}, k)$. There are two indices involved in this construction, t and k apart from the one appearing when taking homotopy groups, but thanks to the suspension theorem 2.12, one index amongst t and k is redundant as

$$M_{\bullet}(X, \mathbb{P}^t/\mathbb{P}^{t-1}, k) \xrightarrow{\sim} M_{\bullet}(X, \mathbb{P}^{t+1}/\mathbb{P}^t, k + 1)$$

is a homotopy equivalence. What really matters is the difference $q = t - k$. So we get rid of the useless index in a way that ensures good functorial properties in situations where the suspension is involved.

Definition 2.14. The *morphic space* of level q associated to a quasi-projective algebraic variety X is the simplicial abelian group $M_{\bullet}(X, q)$ given as the colimit

$$M_{\bullet}(X, q) = \text{colim}_k M_{\bullet}(X, \mathbb{P}^{q+k}/\mathbb{P}^{q+k-1}, k). \quad (11)$$

where the maps

$$M_{\bullet}(X, \mathbb{P}^{q+k}/\mathbb{P}^{q+k-1}, k) \xrightarrow{\sim} M_{\bullet}(X, \mathbb{P}^{q+k+1}/\mathbb{P}^{q+k}, k + 1)$$

are the homotopy equivalences in the suspension theorem.

The *morphic cohomology groups* of X are the groups $L^q H^n(X)$ given by

$$L^q H^n(X) = \pi_{2q-n} M_{\bullet}(X, q). \quad (12)$$

Remark 2.15. All the maps $M_\bullet(X, \mathbb{P}^{q+k}/\mathbb{P}^{q+k-1}, k) \rightarrow M_\bullet(X, q)$ are homotopy equivalences. In particular,

$$L^q H^n(X) \simeq \pi_{2q-n} M_\bullet(X, \mathbb{P}^{q+k}/\mathbb{P}^{q+k-1}, k) \quad (13)$$

for any $k \geq 0$.

Recall that, for an abelian category $\mathcal{A}$, the Dold-Kan correspondence establishes an equivalence

$$N: s\mathcal{A} \rightarrow \mathbf{Ch}_+(\mathcal{A}) \quad (14)$$

between the categories $s\mathcal{A}$ of simplicial objects in $\mathcal{A}$ and the category $\mathbf{Ch}_+(\mathcal{A})$ of positively graded chain complexes in $\mathcal{A}$. For every simplicial object $A_\bullet$ in $s\mathcal{A}$, the associated normalized chain complex $N_*(A)$ is such that

$$\pi_n A_\bullet \simeq H_n N_*(A). \quad (15)$$

Applying the Dold-Kan functor N to the morphic cohomology spaces $M_\bullet(X, q)$ we obtain a chain complex of abelian groups. It is convenient, in analogy with the motivic world, to reindex those morphic complexes as follows:

$$M^n(X, q) = M_{2q-n}(X, q), \quad (16)$$

obtaining a cochain complex of abelian groups $M^*(X, q)$, which is zero for $n > 2q$, so it is bounded above, but is unbounded below.

There is a couple of important properties we will use.

Theorem 2.16. *Let X be an algebraic variety and $p: E \rightarrow X$ be a vector bundle over X . Then the induced map*

$$M_\bullet(X, q) \rightarrow M_\bullet(E, q) \quad (17)$$

is a weak homotopy equivalence and, in particular, the respective morphic cohomology groups are isomorphic.

Proof. Proposition 3.5 in [Fri98] tells us that the map $M_\bullet(X, Y, q) \rightarrow M_\bullet(E, Y, q)$ between bivariant spaces is a homotopy equivalence for Y projective. Then a five lemma argument applied to the pair $\mathbb{P}^q/\mathbb{P}^{q-1}$ proves what we want. $\bullet$

Theorem 2.17 (Duality). *Let X be a quasi-projective variety of dimension d . There is a natural map*

$$\Gamma: M_\bullet(X, q) \rightarrow Z_\bullet(X \times \mathbb{A}^q, d) \quad (18)$$

called the graphing construction.

Moreover, when X is smooth, Γ becomes a homotopy equivalence. In such case gives isomorphisms

$$L^q H^n(X) \xrightarrow{\sim} L_{d-q} H_{2d-n}(X). \quad (19)$$

Proof. The map Γ is constructed and proved to be a homotopy equivalence for smooth varieties in [FL97] theorem 3.3 when X is projective and extended to the quasi-projective case in [Fri98] theorem 5.2. The isomorphism (19) comes from the combination of the homotopy equivalence Γ and the homotopy invariance for cycle spaces 2.13. •

Corollary 2.18. *Let X be smooth and $\{U, V\}$ an open covering of X . Then the diagram*

$$\begin{array}{ccc} M^*(X, q) & \longrightarrow & M^*(U, q) \\ \downarrow & & \downarrow \\ M^*(V, q) & \longrightarrow & M^*(U \cap V, q) \end{array} \quad (20)$$

gives rise to a Mayer-Vietoris long exact sequence for morphic cohomology

$$\cdots \longrightarrow L^q H^n(X) \longrightarrow L^q H^n(U) \oplus L^q H^n(V) \longrightarrow L^q H^n(U \cap V) \longrightarrow \cdots \quad (21)$$

Proof. As X is smooth, The duality theorem implies that the square (20) is quasi-isomorphic to the diagram obtained applying the Dold-Kan functor to

$$\begin{array}{ccc} Z_\bullet(X \times \mathbb{A}^q, d) & \longrightarrow & Z_\bullet(U \times \mathbb{A}^q, d) \\ \downarrow & & \downarrow \\ Z_\bullet(V \times \mathbb{A}^q, d) & \longrightarrow & Z_\bullet(U \cap V \times \mathbb{A}^q, d) \end{array}$$

That last square is homotopy cartesian, because by the localization theorem 2.7 the homotopy fibers of the horizontal maps are $Z_\bullet((X \setminus U) \times \mathbb{A}^q, d)$ and $Z_\bullet((V \setminus U) \times \mathbb{A}^q, d)$, and those spaces are homotopy equivalent (in fact isomorphic) as $X \setminus U = V \setminus U$. •

2.3 A modification of the morphic cohomology

We are interested in using homotopy invariance and Mayer-Vietoris properties for morphic cohomology. Those properties hold for smooth varieties thanks to duality (theorems 2.16 and 2.18) but unfortunately are not known (at least to the author) to hold for singular varieties. So we modify the definition of morphic cohomology to force those two properties. This is not new. In [Fri00] Friedlander defines a modified morphic cohomology, called topological cycle cohomology, which satisfies a Mayer-Vietoris property for Zariski open covers. In fact, the point of [Fri00] is to prove that this theory together with a homological companion satisfies the Bloch-Ogus axioms.

First of all, note that the contravariant functoriality of the complexes $M^*(X, q)$ with respect to X makes them a cochain complexes of presheaves of abelian groups on the category $\mathbf{qProj}_{\mathbb{C}}$.

Let $\Delta_\bullet$ be the standard cosimplicial scheme

$$\Delta_n = \{(x_0, \dots, x_n) \in \mathbb{A}^{n+1} \mid x_0 + \cdots + x_n = 1\}.$$

Definition 2.19. We define the *morphic complexes of sheaves* $\mathcal{M}_{\text{zar}}^*(q)$ as the Zariski sheafification of the presheaf of complexes $M^*(-, q)$, that is

$$\mathcal{M}_{\text{zar}}^n(q) = \text{sh}_{\text{zar}} M^*(-, q). \quad (22)$$

Additionally, we define the *homotopy invariant morphic complexes* as the following total complex

$$\mathcal{M}_{\text{hi}}^n(q) = \text{Tot}^n \mathcal{M}_{\text{zar}}^*(- \times \Delta_{\bullet}, q). \quad (23)$$

Remark 2.20. The complexes $\mathcal{M}^*(q)$ are unbounded below and bounded above by $2q$. This poses some homological algebra troubles as they are bounded on the wrong side. However, due to a result of Spaltenstein [Spa88] one can still have resolutions $\mathcal{M}^*(q) \rightarrow \mathcal{I}^*$ playing the role of injective resolutions. Those are called K-injective in [Spa88]. Later, we will use cohomological finiteness arguments to prove convergence of the spectral sequences we encounter.

We will think of the complexes $\mathcal{M}_{\text{zar}}^*(q)$ and $\mathcal{M}_{\text{hi}}^*(q)$ as objects in the derived category $\mathbf{D}^- \mathbf{Sh}(\mathbf{qProj}_{\mathbb{C}})$ of abelian sheaves on the Zariski site $\mathbf{qProj}_{\mathbb{C}}$.

For any variety X , and a complex of sheaves $\mathcal{F}^* \in \mathbf{D}^- \mathbf{Sh}(\mathbf{qProj}_{\mathbb{C}})$, we will denote by $\mathcal{F}^*|_X$ the restriction of $\mathcal{F}^*$ to the small zariski site of X , i.e. $\mathcal{F}^*|_X$ is an element in the category $\mathbf{D}^- \mathbf{Sh}(X)$, the derived category of abelian sheaves on X .

This sheaf-theoretic interpretation leads to natural reformulations of the morphic cohomology groups.

Definition 2.21. The *topological cycle cohomology groups* are the hypercohomology groups

$$L^q H_{\text{zar}}^n(X) = \mathbb{H}^n(X, \mathcal{M}_{\text{zar}}(q)|_X). \quad (24)$$

The *homotopy invariant morphic cohomology* are the hypercohomology groups

$$L^q H_{\text{hi}}^n(X) = \mathbb{H}^n(X, \mathcal{M}_{\text{hi}}(q)|_X). \quad (25)$$

Remark 2.22. The topological cycle cohomology groups $L^q H_{\text{zar}}^n(X)$ are the ones defined by Friedlander in [Fri00]. Although the original morphic cohomology groups are homotopy invariant by 2.16, we are not able to prove this homotopy invariance for $L^q H_{\text{zar}}^n$. This is the reason for introducing the theory $L^q H_{\text{hi}}^n$ which exhibits both properties, homotopy invariance and Mayer-Vietoris.

Remark 2.23. The hypercohomologies in 2.21 can be rephrased as an ext-group on the derived category $\mathbf{D}^- \mathbf{Sh}(X)$ of sheaves of abelian groups on X , for example

$$L^q H_{\text{hi}}^n(X) = \text{Ext}_{\mathbf{Sh}(X)}^n(\mathbb{Z}_X, \mathcal{M}_{\text{hi}}^*(q)|_X), \quad (26)$$

where $\mathbb{Z}_X$ denotes the constant sheaf on X .

Moreover, let $\mathbb{Z}h_X$ denote the free abelian group generated by the sheaf on $\mathbf{qProj}_{\mathbb{C}}$ represented by X , i.e. $\mathbb{Z}h_X(U) = \mathbb{Z}\mathrm{Hom}(U, X)$. Then there is an isomorphism

$$L^q H_{\mathrm{hi}}^n(X) = \mathrm{Ext}_{\mathbf{Sh}(\mathbf{qProj}_{\mathbb{C}})}^n(\mathbb{Z}h_X, \mathcal{M}_{\mathrm{hi}}^*(q)), \quad (27)$$

where now the Ext is taken in the category of sheaves on the big site $\mathbf{qProj}_{\mathbb{C}}$.

We will use either interpretation, as an Ext in the big site or in the small site.

This new versions of morphic cohomology are related to the old one as follows. There are comparison morphisms

$$M^*(-, q) \xrightarrow{a} \mathcal{M}_{\mathrm{zar}}^*(q) \xrightarrow{b} \mathcal{M}_{\mathrm{hi}}^*(q), \quad (28)$$

the first is the inclusion of a presheaf in its associated sheaf, and the second is the inclusion into the summand of the total complex corresponding to the algebraic simplex Δ_0 .

As defined in 2.14, $L^q H^n(X)$ is the homology of the complex of global sections of the cochain complex of presheaves $M^*(-, q)$, while the other two flavours of morphic cohomology are the hypercohomology of the respective sheaves.

Let's take then K-injective resolutions $\mathcal{M}_{\mathrm{zar}}^*(q) \rightarrow \mathcal{I}_{\mathrm{zar}}^*$ and $\mathcal{M}_{\mathrm{hi}}^*(q) \rightarrow \mathcal{I}_{\mathrm{hi}}^*$. There are induced comparison maps

$$M^*(-, q) \xrightarrow{a} \mathcal{I}_{\mathrm{zar}}^* \xrightarrow{b} \mathcal{I}_{\mathrm{hi}}^*,$$

which, when taking cohomology of the global sections induce comparison maps

$$L^q H^n(X) \xrightarrow{a} L^q H_{\mathrm{zar}}^n(X) \xrightarrow{b} L^q H_{\mathrm{hi}}^n(X). \quad (29)$$

Proposition 2.24. *Assume X is a smooth quasi-projective variety. Then the comparison morphisms (29) are isomorphisms.*

Proof. From 2.18 we know that the presheaf $M_{\bullet}(-, q)$ restricted to smooth varieties satisfies the Mayer-Vietoris property, that is, the square

$$\begin{array}{ccc} M^*(X, q) & \longrightarrow & M^*(U, q) \\ \downarrow & & \downarrow \\ M^*(V, q) & \longrightarrow & M^*(U \cap V, q) \end{array}$$

is homotopy cartesian. Take a K-injective resolution $\mathcal{M}_{\mathrm{zar}}^*(q) \rightarrow \mathcal{I}_{\mathrm{zar}}^*$. Using the Brown-Gersten theorem ([Voe00] lemma 3.5) we conclude that the presheaf $M^*(-, q)$ is globally weakly equivalent to the K-injective resolution $\mathcal{I}^*$, so

$$H^n \Gamma(X, M^*(-, q)) \simeq H^n \Gamma(X, \mathcal{I}_{\mathrm{zar}}^*),$$

and this settles the first isomorphism a .

As for b , by theorem 2.16 we know that $L^q H^n(-)$ is a homotopy invariant functor, and by the previous isomorphism coincides with $L^q H_{\text{zar}}^n(-)$ on smooth varieties, so the last one is also homotopy invariant on smooth varieties. Then the spectral sequence associated to the double complex defining $L^q H_{\text{hi}}^n$ degenerates on the second page and this gives the isomorphism b . ●

Finally, those are the properties we will use of the modified theory $L^q H_{\text{hi}}^n$.

Theorem 2.25. *Let U, V be an open cover of a complex variety X . Then there is a Mayer-Vietoris long exact sequence for the groups $L^q H_{\text{hi}}^n$*

$$\cdots \longrightarrow L^q H_{\text{hi}}^n(X) \longrightarrow L^q H_{\text{hi}}^n(U) \oplus L^q H_{\text{hi}}^n(V) \longrightarrow L^q H_{\text{hi}}^n(U \cap V) \longrightarrow \cdots \quad (30)$$

Proof. The open cover $\{U, V\}$ gives an exact triangle in $\mathbf{D}^- \mathbf{Sh}(X)$

$$\mathbb{Z}_{U \cap V} \longrightarrow \mathbb{Z}_U \oplus \mathbb{Z}_V \longrightarrow \mathbb{Z}_X \longrightarrow \mathbb{Z}_{U \cap V}[1] \quad (31)$$

which, when passed through the functor $\text{Ext}^n(-, \mathcal{M}_{\text{hi}}^*(q))$, gives the desired long exact sequence. ●

Theorem 2.26. *The groups $L^q H^n(X)$ are homotopy invariant, that is the projection $p: X \times \mathbb{A}^1 \rightarrow X$ induces isomorphisms*

$$L^q H^n(X) \xrightarrow{\sim} L^q H^n(X \times \mathbb{A}^1). \quad (32)$$

Proof. This is a standard argument. See for example corollary 2.19 in [MVW06] ●

2.4 Some remarks about notation

Remark 2.27. In the remainder of this paper we will deal only with the version $L^q H_{\text{hi}}^n$ of morphic cohomology, as we are interested in using a version of a Mayer-Vietoris spectral sequence and homotopy invariance. We will drop the subindex “hi” from the notation for readability.

It will be useful to deal with the morphic complexes all at once, so we define the complex of sheaves

$$\mathcal{M}^* = \bigoplus_{q \geq 0} \mathcal{M}^*(q).$$

This is a bigraded object, having the algebraic degree q and the cohomological degree n with a differential for the n -grading. We will denote by $\mathcal{H}^n \mathcal{M}$ its cohomology sheaves.

Analogously, we will use the notation LH^n for the graded group

$$LH^n(X) = \bigoplus_{q \geq 0} L^q H^n(X) = \mathbb{H}^n(X, \mathcal{M}^*),$$

and LH for the bigraded groups

$$LH(X) = \bigoplus_{n \geq 0} LH^n(X).$$

2.5 Cup product and Kunneth homomorphism

Let X, X' be schemes. Following [FW01] proposition 3.2, the projections $\pi_X: X \times X' \rightarrow X$ and $\pi_{X'}: X \times X' \rightarrow X'$ induce exterior products on morphic complexes

$$M^*(X, q) \otimes M^*(X', q') \rightarrow M^*(X \times X', q + q'). \quad (33)$$

Composing this exterior product with the pull-back through the diagonal embedding $\Delta: X \rightarrow X \times X$ we get a cup product

$$M^*(X, q) \otimes M^*(X, q') \rightarrow M^*(X \times X, q + q') \rightarrow M^*(X, q + q'), \quad (34)$$

which induces a cup product

$$\mathcal{M}^*(q) \otimes \mathcal{M}^*(q') \rightarrow \mathcal{M}^*(q + q'). \quad (35)$$

on the sheaves $\mathcal{M}^*$.

Theorem 2.28. *The cup product (35) induces an associative product on morphic cohomology*

$$L^q H^n(X) \otimes L^{q'} H^{n'}(X) \rightarrow L^{q+q'} H^{n+n'}(X). \quad (36)$$

which is graded commutative with respect to the cohomological grading, that is, for $a \in LH^n(X)$, $b \in LH^{n'}(X)$ the commutativity relation

$$a \cup b = (-1)^{nn'} b \cup a$$

holds.

Moreover, the cup product is functorial in the sense that for $f: X \rightarrow Y$ a morphism of algebraic varieties, the induced map on morphic cohomology $f^*: LH(Y) \rightarrow LH(X)$ is a ring homomorphism.

Proof. See [FL92] corollary 6.2. ●

Let $\mathcal{M}^*(\text{pt}) = \Gamma(\text{Spec } \mathbb{C}, \mathcal{M}^*)$ be the sections of the morphic complex on a point, and let $LH(\text{pt})$ be its hypercohomology, which in this case coincides with the cohomology ring of the global sections $\mathcal{M}^*(\text{pt})$.

Corollary 2.29. *The morphic cohomology of a complex quasi-projective variety X has a canonical structure of $LH(\text{pt})$ -module.*

Proof. Applying the second part of theorem 2.28 to the structure map $X \rightarrow \text{Spec } \mathbb{C}$ we get a ring homomorphism $LH(\text{pt}) \rightarrow LH(X)$, making $LH(X)$ into a $LH(\text{pt})$ -module. ●

Now, the exterior product (33) also induces an exterior product on morphic cohomology

$$L^q H^n(X) \otimes L^{q'} H^{n'}(X') \rightarrow L^{q+q'} H^{n+n'}(X \times X'). \quad (37)$$

As the structure of $LH(\text{pt})$ -module in $LH(X)$ is functorial, the action of $LH(\text{pt})$ on either factor on the left of (37), gives the same action on the right, so the exterior product map (37) factors through

$$LH(X) \otimes_{LH(\text{pt})} LH(Y) \rightarrow LH(X \times Y). \quad (38)$$

In rather special circumstances this Kunneth homomorphism (38) happens to be an isomorphism. We are interested in a very special case of this Kunneth isomorphism, which we now prove.

Proposition 2.30. *Let X be a smooth quasi-projective variety. The Künneth homomorphism*

$$LH(X) \otimes_{LH(\text{pt})} LH(\mathbb{G}_m) \rightarrow LH(X \times \mathbb{G}_m). \quad (39)$$

is an isomorphism.

Proof. Let $i: \text{pt} \rightarrow \mathbb{A}^1$ be the inclusion of a point and $j: \mathbb{G}_m \rightarrow \mathbb{A}^1$ its open complement. We have the following commutative diagram of long exact sequences

$$\begin{array}{ccccccc} \cdots LH(X) \otimes_{LH(\text{pt})} LH(\text{pt}) & \xrightarrow{id \times i_!} & LH(X) \otimes_{LH(\text{pt})} LH(\mathbb{A}^1) & \xrightarrow{id \times j^*} & LH(X) \otimes_{LH(\text{pt})} LH(\mathbb{G}_m) & \cdots \\ \downarrow & & \downarrow & & \downarrow & \\ \cdots LH(X \times \text{pt}) & \xrightarrow{(id \times i)_!} & LH(X \times \mathbb{A}^1) & \xrightarrow{(id \times j)^*} & LH(X \times \mathbb{G}_m) & \cdots \end{array}$$

The vertical maps are the Kunneth morphisms, and $i_!$ is the Gysin map defined by duality (theorem 2.17) as $i_! = \Gamma_*^{-1} i_* \Gamma_*$. The exactness of the rows come, by duality, from the localization theorem 2.7 and the long exact sequence of homotopy groups of a fibration.

Now a standard application of the five lemma together with the homotopy invariance 2.16 proves the desired isomorphism. ●

Remark 2.31. I became aware of a construction of a Kunneth spectral sequence in the case in which X or Y is a linear variety in a private communication with Mircea Voineagu [Voi08a]. An analogous construction is done in [Jos01] for higher Chow groups and K-theory.

Finally, some basic computations of morphic cohomology rings we will need.

Proposition 2.32. 1. For $k \geq 0$,

$$LH(\mathbb{A}^k) \simeq \mathbb{Z}[s],$$

where s is a free generator of bidegree $(1, 0)$ (degree 1 with respect to the q -grading).

2. For $k \geq 0$,

$$LH(\mathbb{P}^k) \simeq \mathbb{Z}[s, h]/(h^{k+1})$$

where s has bidegree $(1, 0)$ and h has bidegree $(1, 2)$.

3. The morphic cohomology of the multiplicative group is given by

$$LH(\mathbb{G}_m) \simeq \mathbb{Z}[s, e]/(e^2),$$

where s is a generator of bidegree $(1, 0)$ and e is a generator of bidegree $(1, 1)$.

Proof. 1) and 2) follow from duality and the computations of Lawson homology using the suspension theorem 2.12.

3) Take the open cover of $\mathbb{P}^1$ by two affines. Then we have the following piece of Mayer-Vietoris sequence

$$LH^n(\mathbb{P}^1) \longrightarrow LH^n(\mathbb{A}^1)^{\oplus 2} \longrightarrow LH^n(\mathbb{G}_m) \longrightarrow LH^{n+1}(\mathbb{P}^1) \longrightarrow LH^{n+1}(\mathbb{A}^1)^{\oplus 2}$$

which, using 1) and 2) for the computations of $\mathbb{P}^1$ and $\mathbb{A}^1$ gives the result. ●

Remark 2.33. Note that, in particular, $LH(\text{pt}) \simeq \mathbb{Z}[s]$. As a consequence of 2.29, the morphic cohomology ring $LH(X)$ is, in fact, a $\mathbb{Z}[s]$ -module. The action by s on $LH(X)$ corresponds to the s -maps defined in [FM94].

3 Toric varieties

First we set the notation. Let $N \simeq \mathbb{Z}^n$ be a free $\mathbb{Z}$ -module of rank n , and M its dual $\mathbb{Z}$ -module. We will denote by $N_{\mathbb{R}} = N \otimes \mathbb{R}$ and $M_{\mathbb{R}} = M \otimes \mathbb{R}$. In this way, N and M are to be thought as lattices on $N_{\mathbb{R}}$ and $M_{\mathbb{R}}$. Moreover, there is the duality pairing $\langle u, v \rangle$ for $u \in N_{\mathbb{R}}$ and $v \in M_{\mathbb{R}}$.

3.1 Cones and fans

Definition 3.1. A *rational polyhedral cone* in N is a set $\sigma \subset N_{\mathbb{R}}$ generated by a finite number of integral vectors $v_1, \dots, v_k \in N$ in the following way:

$$\sigma = \{\lambda_1 v_1 + \dots + \lambda_k v_k \in N_{\mathbb{R}} \mid \lambda_i \in \mathbb{R}_{\geq 0}\}.$$

The *dimension* of a cone σ is the dimension of the vector space $\mathbb{R}\sigma$ generated by the cone.

If a rational polyhedral cone σ does not contain a vector space of positive dimension is said to be *strictly convex*.

Definition 3.2. Let σ be a rational polyhedral cone in N .

1. The *dual cone* of σ is a rational polyhedral cone in M given by

$$\sigma^\vee = \{v \in M_{\mathbb{R}} \mid \langle u, v \rangle \geq 0 \text{ for all } u \in \sigma\}.$$

2. The *orthogonal cone* of σ is a rational polyhedral cone in M given by

$$\sigma^\perp = \{v \in M_{\mathbb{R}} \mid \langle u, v \rangle = 0 \text{ for all } u \in \sigma\}.$$

Remark 3.3. The orthogonal cone $\sigma^\perp$ is in fact a vector space, as if $\langle u, v \rangle = 0$ then $\langle u, \lambda v \rangle = 0$ for all $\lambda \in \mathbb{R}$.

Definition 3.4. A cone τ is a *face* of σ if there exists a $u \in M_{\mathbb{R}}$ such that

$$\tau = \sigma \cap u^\perp.$$

Definition 3.5. A *fan* Δ is a set of strongly convex rational polyhedral cones in N such that

1. Every face of a cone in Δ also belongs to Δ .
2. The intersection of two cones in Δ is a face of both intersecting cones.

The notation $\Delta^{(k)}$ will mean the set of all cones of codimension k in Δ .

Remark 3.6. A fan has a partial order given by the inclusion of faces. We will use the notation $\tau \leq \sigma$ to say that τ is a face of σ .

Definition 3.7. An *orientation of a cone* σ is an orientation of the vector spaces $\mathbb{R}\sigma$.

An *orientation of a fan* Δ will be a choice of an orientation for every cone in Δ .

We will always use fans with a fixed orientation.

Remark 3.8. Let $\tau \leq \sigma$ be a face of codimension 1 in σ . This means that there exists a $u \in \sigma^\vee$ such that $\tau = \sigma \cap u^\perp$. Then we have an isomorphism

$$\mathbb{R}\sigma \simeq \mathbb{R}u \oplus \mathbb{R}\tau. \tag{40}$$

which allows us to transfer the orientation of σ to τ as follows: the orientation induced on τ by σ is the one compatible with the isomorphism (40) and taking the orientation in $\mathbb{R}u$ given by the vector u .

3.2 Definition of a toric variety

Let us fix a field k . For now it will be an arbitrary field, but in the next sections we will set $k = \mathbb{C}$.

We describe the toric variety $X(\Delta)$ associated to a fan Δ . The scheme $X(\Delta)$ is constructed locally, one affine piece for every cone in Δ , and then glueing them, according to the combinatorics of the fan, to build the scheme $X(\Delta)$.

For every cone $\sigma \in \Delta$ we define an affine scheme

$$X_\sigma = \operatorname{Spec} k[\sigma^\vee \cap M],$$

where $\sigma^\vee \cap M$ is seen as a monoid and $k[\sigma^\vee \cap M]$ is its monoid algebra. An inclusion of cones $\tau \leq \sigma$ induces an inclusion on dual cones $\sigma^\vee \leq \tau^\vee$ and this induces a morphism on the corresponding monoid algebras, obtaining a morphism of schemes $i_{\tau,\sigma}: X_\tau \rightarrow X_\sigma$.

Proposition 3.9. *Let Δ be a fan on a lattice N of dimension n .*

1. *For every face inclusion $\tau \leq \sigma$, the induced morphism $i_{\tau,\sigma}: X_\tau \rightarrow X_\sigma$ is an open embedding.*
2. *There is a (unique) k -scheme $X(\Delta)$ together with a family of open embeddings $i_\sigma: X_\sigma \rightarrow X(\Delta)$ for every cone in Δ such that X_σ is an open covering of $X(\Delta)$ and for every pair of cones $\tau \leq \sigma$, the following diagram commutes*

$$\begin{array}{ccc} X_\tau & \xrightarrow{i_{\tau,\sigma}} & X_\sigma \\ & \searrow i_\tau & \downarrow i_\sigma \\ & & X(\Delta) \end{array}$$

3. *The scheme $X(\Delta)$ is reduced, normal and Noetherian. Moreover, if the cones in Δ cover the whole $N_{\mathbb{R}}$ the variety $X(\Delta)$ is complete.*
4. *For $\tau, \sigma \in \Delta$, the face inclusions of $\tau \cap \sigma$ induce an isomorphism*

$$X_{\tau \cap \sigma} \simeq X_\tau \times_{X(\Delta)} X_\sigma.$$

Proof. See [Ful93], sections 1.4 and 2.1. •

Definition 3.10. A *toric variety* for a fan Δ is an algebraic variety $X(\Delta)$ associated Δ as in the previous theorem.

Example 3.11. Let N be a lattice of rank n and Δ the fan containing only the cone $\{0\}$. Then $X(\Delta)$ is an algebraic torus of dimension n , that is there is an isomorphism

$$X(\Delta) = \operatorname{Spec} k[M] \simeq \mathbb{G}_m^n.$$

This algebraic torus is, in fact, a commutative group scheme, and the group law is described by the comultiplication on the coordinate ring $k[M] \rightarrow k[M] \otimes k[M]$ given by $m \mapsto m \otimes m$ for every $m \in M$ and extended by linearity.

Remark 3.12. Every toric variety $X(\Delta)$ contains an algebraic torus as a dense open set, corresponding to the affine piece

$$X_0 = \operatorname{Spec} k[M] \simeq \mathbb{G}_m^n$$

associated to the cone $0 \in \Delta$.

The group law on the torus X_0 can be extended to an action on the whole toric variety. Locally on an open piece $X_\sigma = \operatorname{Spec} k[\sigma^\vee \cap M]$ this action is given by the morphism of rings

$$k[\sigma^\vee \cap M] \rightarrow k[M] \otimes k[\sigma^\vee \cap M]$$

such that $m \mapsto m \otimes m$ for every $m \in \sigma^\vee \cap M$. Then all those local actions glue together forming an action

$$X_0 \times X(\Delta) \rightarrow X(\Delta).$$

To every cone $\sigma \in \Delta$, we can associate an affine scheme T_σ besides the open affine X_σ we have already constructed,

$$T_\sigma = \operatorname{Spec} k[\sigma^\perp \cap M].$$

Observe that There is a homomorphism of rings

$$k[\sigma^\vee \cap M] \rightarrow k[\sigma^\perp \cap M]$$

sending every element of $\sigma^\perp$ to themselves and the elements of $\sigma^\vee$ not in $\sigma^\perp$ to 0. This homomorphism induces a morphism of schemes

$$T_\sigma = \operatorname{Spec} k[\sigma^\perp \cap M] \rightarrow X_\sigma = \operatorname{Spec} k[\sigma^\vee \cap M],$$

and composing with the embedding $i_\sigma: X_\sigma \rightarrow X(\Delta)$ we get morphisms $j_\sigma: T_\sigma \rightarrow X(\Delta)$.

On the other hand, we have retractions $r_\sigma: X_\sigma \rightarrow T_\sigma$ defined by the inclusion $\sigma^\perp \rightarrow \sigma^\vee$.

Proposition 3.13. *For every cone $\sigma \in \Delta$, the morphism*

$$j_\sigma: T_\sigma \rightarrow X(\Delta),$$

is a locally closed embedding and corresponds with an orbit of the torus action.

Moreover, the collection of all T_σ for all the cones $\sigma \in \Delta$ give the orbit decomposition of the torus action on the toric variety $X(\Delta)$.

Proof. See [Ful93] section 3.1. ●

Remark 3.14. Note that the orbits T_σ are algebraic tori whose dimension is the codimension of σ , i.e.

$$\dim T_\sigma = n - \dim \sigma.$$

In short, associated to a toric variety we have an open cover given by the affine pieces

$$X_\sigma = \operatorname{Spec} k[\sigma^\vee \cap M],$$

and an orbit decomposition as a disjoint union of tori

$$T_\sigma = \operatorname{Spec} k[\sigma^\perp \cap M].$$

There is a relation between them in terms of the combinatorics of the fan as follows:

Proposition 3.15. *We have the following disjoint union decompositions,*

$$X_\sigma = \bigcup_{\tau \leq \sigma} T_\tau,$$

$$\overline{T}_\tau = \bigcup_{\tau \leq \sigma} T_\sigma.$$

Proof. See [Ful93] section 3.1. ●

Remark 3.16. In particular, the orbit T_σ is closed in X_σ , and is the lowest dimensional orbit contained in it.

Proposition 3.17. *Let $X(\Delta)$ be a toric variety and σ a cone in Δ . There is a morphism*

$$h: X_\sigma \times \mathbb{A}_k^1 \rightarrow X_\sigma$$

such that

1. $h(-, 1) = \operatorname{id}$,
2. $h(-, 0) = j_\sigma r_\sigma$,
3. $h(-, t)$ restricts to the identity on T_σ for every t .

So, the morphism h gives an algebraic homotopy equivalence between X_σ and T_σ .

Proof. We have $X_\sigma = \operatorname{Spec} k[\sigma^\vee \cap M]$, $T_\sigma = \operatorname{Spec} k[\sigma^\perp \cap M]$. The inclusion $T_\sigma \rightarrow X_\sigma$ is given by the quotient

$$k[\sigma^\vee \cap M] \rightarrow k[\sigma^\perp \cap M],$$

which is the identity on $\sigma^\perp$ and sends any element $v \in \sigma^\vee$ not in $\sigma^\perp$ to $0 \in k[\sigma^\perp \cap M]$.

Pick $u_0 \in \sigma$ such that $\sigma^\perp = \sigma^\vee \cap u_0^\perp$. Then define

$$h^*: k[\sigma^\vee \cap M] \rightarrow k[\sigma^\vee \cap M] \otimes k[t],$$

by $h^*(v) = v \otimes t^{\langle u_0, v \rangle}$ for every $v \in \sigma^\vee$. This gives a morphism of schemes $h: X_\sigma \times \mathbb{A}_k^1 \rightarrow X_\sigma$ with the desired properties. $\bullet$

3.3 Morp hic cohomology of an algebraic torus

Now we compute the morp hic cohomology ring of an algebraic torus. As we will need this computation for subtori of a toric variety, it will be useful to have a canonical description of this ring in terms of the lattice defining the toric variety.

Let N be a lattice of rank n , and $L_\mathbb{R} \subset M_\mathbb{R} = N_\mathbb{R}^\vee$ a subspace of dimension r generated by vectors in the lattice M . Consider the rank r sublattice $L = L_\mathbb{R} \cap M$, and its associated torus

$$T_L = \operatorname{Spec} \mathbb{C}[L].$$

As we have seen in 2.32, $LH^1(\mathbb{G}_m) \simeq \mathbb{Z}[s]e$ with e a generator in bidegree $(1,1)$ which corresponds, by duality, to a radial Borel-Moore chain joining 0 and ∞ in $\mathbb{G}_m$. Now, any $v \in L$ defines a character $\chi_v: T_L \rightarrow \operatorname{Spec} \mathbb{C}[v, v^{-1}] = \mathbb{G}_m$, so we can now define a morphism of rings

$$\varphi: (\wedge^* L) \otimes LHpt \rightarrow LH^*(T_L) \tag{41}$$

by

$$\varphi(v \otimes 1) = \chi_v^*(e),$$

for $v \in L$, and extended in the obvious way to the exterior algebra because $LH^*(T_L)$ is a graded commutative algebra (theorem 2.28). On the left hand side of (41) the vectors in the lattice L are assumed to have bigrading $(1,1)$.

Theorem 3.18. *The morphism φ is an isomorphism.*

Proof. We argue by induction on the rank of L . The isomorphism is clear when $\operatorname{rank} L = 1$ by the computation in 2.32. Let $L = L_0 \oplus \mathbb{Z}v$. This gives a product decomposition $T_L =$

$T_{L_0} \times \mathbb{G}_m$. Now, because the Kunneth isomorphism in 2.30 preserves the cup product, we get a commutative diagram

$$\begin{array}{ccc} \wedge^n L \otimes LH(\text{pt}) & \longrightarrow & \bigoplus_{p+q=n} (\wedge^p L_0 \otimes LH(\text{pt})) \otimes_{LH(\text{pt})} (\wedge^q \mathbb{Z}v \otimes LH(\text{pt})) \\ \downarrow \varphi & & \downarrow \\ LH^n(T_L) & \longrightarrow & \bigoplus_{p+q=n} LH^p(T_{L_0}) \otimes_{LH(\text{pt})} LH^q(\mathbb{G}_m) \end{array}$$

The upper row is an isomorphism by multilinear algebra results, while the lower row is an isomorphism by the Kunneth isomorphism 2.30. The right vertical map is a sum of tensor products of φ 's corresponding to lower dimensional tori, so are isomorphisms by induction hypothesis. We conclude then that the left vertical map is an isomorphism. ●

4 Spectral sequence associated to a toric variety

Let $X(\Delta)$ be a complete toric variety of dimension n , R a ring (possibly graded) and $\mathcal{F}^*$ a cochain complex of sheaves of R -modules on X . As usual, the hypercohomology of $\mathcal{F}^*$ will be

$$\mathbb{H}^n(X(\Delta), \mathcal{F}^*) = H^n \Gamma(X(\Delta), \mathcal{I}^*)$$

where $\mathcal{I}^*$ is a K-injectives resolution $\mathcal{F}^* \rightarrow \mathcal{I}^*$.

In this section we will write down a spectral sequence converging to the hypercohomology $\mathbb{H}^n(X(\Delta), \mathcal{F}^*)$ whose E_2 page is computable in terms of the combinatorics of the toric variety, and the hypercohomology of $\mathcal{F}^*$ on algebraic tori. The spectral sequence comes from the identification

$$\mathbb{H}^n(X(\Delta), \mathcal{F}^*) = \text{Ext}^n(R_X, \mathcal{F}^*),$$

and the fact that the hyper-ext can be computed resolving either variable. We will chose to resolve the constant sheaf R_X producing a Čech-like resolution $\check{C}_*(\Delta, R_X) \rightarrow R_X$ from the combinatorics of the toric variety.

A similar idea to ours, applied to singular homology and cohomology, was previously developed in the thesis [Jor98].

4.1 Resolution associated to a fan

Let $X(\Delta)$ be a complete toric variety defined by a fan Δ , and let R be a commutative ring.

Definition 4.1. Let $\check{C}_k(\Delta, R)$ for $k \geq 0$ be the sequence of free R -modules

$$\check{C}_k(\Delta, R) = \bigoplus_{\sigma \in \Delta^{(k)}} R[\sigma],$$

with basis $\Delta^{(k)}$, together with R -morphisms $d_k: \check{C}_k(\Delta, R) \rightarrow \check{C}_{k-1}(\Delta, R)$ defined by

$$d_k([\tau])_\sigma = \epsilon(\tau, \sigma),$$

where $\epsilon(\tau, \sigma) = \pm 1$ according to whether the orientation induced by σ on τ coincides or not with the orientation in τ (remember, we assume a fixed choice of orientation on every cone).

Definition 4.2. Let $\check{C}_k(\Delta, R_X)$ for $k \geq 0$ be the sequence of sheaves of R -modules on $X(\Delta)$ given by

$$\check{C}_k(\Delta, R_X) = \bigoplus_{\sigma \in \Delta^{(k)}} i_{\sigma!} i_{\sigma}^* R_X,$$

where R_X is the constant sheaf on $X(\Delta)$ and $i_{\sigma}: X_{\sigma} \rightarrow X(\Delta)$ is the inclusion of the open X_{σ} .

Moreover, we define, a sequence of morphisms $d_k: \check{C}_k(\Delta, R_X) \rightarrow \check{C}_{k-1}(\Delta, R_X)$ given by

$$d_k = \bigoplus_{\substack{\sigma \in \Delta^{(k-1)} \\ \tau \in \Delta^{(k)} \\ \tau \leq \sigma}} \epsilon(\tau, \sigma) \iota_{\tau, \sigma}$$

where $\iota_{\tau, \sigma}: i_{\tau!} i_{\tau}^* R_X \rightarrow i_{\sigma!} i_{\sigma}^* R_X$ is the natural inclusion of sheaves inducing the identity on the nonzero fibers.

Definition 4.3. Given a fan Δ and a cone $\sigma \in \Delta$ of codimension k , we define a fan Δ_{σ} defined on the lattice $N/(\mathbb{R}\sigma \cap N)$ of dimension k , whose cones are the projection of cones in Δ having σ as a face.

Remark 4.4. Notice that the cones in Δ_{σ} correspond bijectively with the cones $\tau \in \Delta$ having σ as a face.

Let $x \in X(\Delta)$ be a point. We denote by $\sigma(x)$ the unique cone in Δ such that $x \in T_{\sigma(x)}$.

Remark 4.5. Observe that

$$(i_{\tau!} i_{\tau}^* R_X)_x = \begin{cases} R & \text{if } \sigma(x) \leq \tau, \\ 0 & \text{otherwise.} \end{cases}$$

In other words, the fiber $(i_{\tau!} i_{\tau}^* R_X)_x$ is nonzero exactly for the cones $\tau \in \Delta_{\sigma(x)}$.

This allows us to define a morphism of R -modules

$$f_{k,x}: \check{C}_k(\Delta, R_X)_x \rightarrow \check{C}_k(\Delta_{\sigma(x)}, R)$$

which is the identity on every nonzero summand, as the summands on the right correspond exactly to the nonzero summands on the left.

Proposition 4.6. *Let $X(\Delta)$ be a complete toric variety associated to a fan Δ .*

1. *The sequence of R -modules $\check{C}_k(\Delta, R)$ together with the differentials d_k form a chain complex of R -modules, that is,*

$$d_{k-1}d_k = 0.$$

2. *The diagram*

$$\begin{array}{ccc} \check{C}_k(\Delta, R_X)_x & \xrightarrow{f_{k,x}} & \check{C}_k(\Delta_{\sigma(x)}, R) \\ \downarrow d_k & & \downarrow d_k \\ \check{C}_{k-1}(\Delta, R_X)_x & \xrightarrow{f_{k-1,x}} & \check{C}_{k-1}(\Delta_{\sigma(x)}, R) \end{array}$$

is commutative, and the rows are isomorphisms.

3. *The sequence of sheaves of R -modules $\check{C}_k(\Delta, R_X)$ together with the differentials d_k form a chain complex of sheaves of R -modules.*

Proof. 1) Let $[\tau]$ be an element of the basis of $\check{C}_k(\Delta, R)$. Then we have

$$d_{k-1}d_k([\tau])_\eta = \sum_{\eta < \sigma < \tau} \epsilon(\eta, \sigma)\epsilon(\sigma, \tau)$$

and either there is no σ in between η and τ or there are exactly two of them, giving opposite signs.

2) Follows easily from the definition and remark 4.5.

3) From 1 and 2 together we see that $d_{k-1,x}d_{k,x}: \check{C}_k(\Delta, R_X)_x \rightarrow \check{C}_{k-2}(\Delta, R_X)_x$ is zero. Then as $(d_{k-1}d_k)_x = d_{k-1,x}d_{k,x}$ and a morphism of sheaves which is zero on the fibers is the zero morphism, we are done. $\bullet$

Definition 4.7. Let $a: \check{C}_*(\Delta, R) \rightarrow R$ be the augmentation morphism given by

$$a(x) = \sum_{\tau \in \Delta^{(0)}} x_\tau, \quad (42)$$

where $x = \sum x_\tau[\tau] \in \check{C}_0(\Delta, R)$.

In a similar way, let $a: \check{C}_*(\Delta, R_X) \rightarrow R_X$ be the augmentation morphism induced by the morphisms $i_{\sigma!}i_{\sigma}^*R_X \rightarrow R_{X(\Delta)}$.

We will prove that $a: \check{C}_*(\Delta, R_X) \rightarrow R_X$ is a quasi-isomorphism. To do so, we will relate the fiber complexes $\check{C}_*(\Delta, R_X)_x$ with the cellular homology complex of a cellular decomposition on the n -dimensional ball.

Let

$$B = \{x \in N_{\mathbb{R}} \mid \|x\| \leq 1\}, \quad (43)$$

$$S = \{x \in N_{\mathbb{R}} \mid \|x\| = 1\}. \quad (44)$$

Definition 4.8. For every non-zero cone $\sigma \in \Delta$ we define the subset $e_\sigma \subset S$ as follows

$$e_\sigma = \sigma \cap S.$$

Proposition 4.9. *The subsets $e_\sigma \subset S$ are homeomorphic to closed balls of dimension $\dim \sigma - 1$. Together form a cellular decomposition of the sphere S . The sphere together with this decomposition will be denoted by S_Δ .*

Proof. As e_σ are the intersection of a strictly convex cone with the unit sphere S_Δ it is clear that are homeomorphic to balls of dimension $\dim \sigma - 1$. They cover all S because the fan is complete. Finally, the boundary of the cell e_σ is formed by the cells associated to the faces of σ , and so belong to the lower dimensional skeleton. $\square$

Definition 4.10. To any complete fan Δ we associate an abstract simplicial complex $K(\Delta)$ as follows:

1. The vertices in $K(\Delta)$ correspond to the cones in Δ .
2. The k -simplexes in $K(\Delta)$ are the sets of vertices belonging to flags in Δ of length k , that is, sequences of strictly included cones

$$\tau_0 < \tau_1 < \cdots < \tau_k.$$

Remark 4.11. The simplicial complex $K(\Delta)$ is related to the barycentric subdivision of the fan Δ . However, they are not exactly the same thing because of the cone $0 \in \Delta$. If we had not admitted the cone 0 in the definition of $K(\Delta)$ we would have obtained a combinatorial model of the barycentric subdivision of the fan Δ .

For every 1-dimensional cone in $\tau \in \Delta^{(n-1)}$ let $u_\tau \in N_{\mathbb{R}}$ be the unique unit vector generating it. Then, for any non-zero cone $\sigma \in \Delta$, let v_σ be the vector

$$v_\sigma = \sum_{\substack{\tau \in \Delta^{(n-1)} \\ \tau \leq \sigma}} u_\tau.$$

Definition 4.12. For every k -simplex $(\tau_0, \dots, \tau_k) \in K(\Delta)$ given by a flag of cones $\tau_0 < \cdots < \tau_k$, we define a subset $d_{(\tau_0, \dots, \tau_k)} \subset B$ as follows,

$$d_{(\tau_0, \dots, \tau_k)} = \begin{cases} \{0\} & \text{if } \tau_0 = 0 \text{ and } k = 0, \\ \mathbb{R}_{\geq 0} \langle v_{\tau_1}, \dots, v_{\tau_k} \rangle \cap B & \text{if } \tau_0 = 0 \text{ and } k > 0, \\ \mathbb{R}_{\geq 0} \langle v_{\tau_0}, \dots, v_{\tau_k} \rangle \cap S & \text{if } \tau_0 \neq 0. \end{cases} \quad (45)$$

Proposition 4.13. *The subsets $d_{\tau_0, \dots, \tau_k} \subset B$ are homeomorphic to closed balls of dimension k . Together form a cellular decomposition of the ball B . The ball together with this decomposition will be denoted by $B_{K(\Delta)}$.*

Proof. Let $(\tau_0, \dots, \tau_k) \in K(\Delta)$. Because the vectors v_{τ_i} all belong to the cone τ_k , the subsets $\mathbb{R}_{\geq 0} \langle v_{\tau_0}, \dots, v_{\tau_k} \rangle$ are strongly convex cones. Then arguing as in 4.9 for every case in definition 4.12 we prove that $d_{\tau_0, \dots, \tau_k}$ are balls. As for the statement regarding its dimension, it follows from the linear independence of the v_{τ_i} for any flag $0 \neq \tau_0 < \dots < \tau_k$.

Finally, observe that the boundary of a cell $d_{\tau_0, \dots, \tau_k}$ is formed by the cells resulting from removing one cone in the flag, all of lower dimension. This proves that the cells $d_{\tau_0, \dots, \tau_k}$ give a cellular decomposition of the ball B . ●

Definition 4.14. For every cone $\sigma \in \Delta$, let $e_\sigma^\vee \subset B$ be the subset defined by

$$e_\sigma^\vee = \bigcup_{\substack{k \geq 0 \\ (\tau_0, \dots, \tau_k) \in K(\Delta) \\ \sigma \leq \tau_0}} d_{\tau_0, \dots, \tau_k} \quad (46)$$

Proposition 4.15. The subsets $e_\sigma^\vee \subset B$ are homeomorphic to closed balls of dimension $\text{codim } \sigma$. Together form a cellular decomposition of the ball B . The ball together with this decomposition will be denoted by $B_\Delta^\vee$.

Proof. Observe that $e_\sigma^\vee$ is a geometric realization of a subcomplex of $K(\Delta)$. Moreover, this subcomplex happens to be isomorphic to $K(\Delta_\sigma)$ (follows directly from the definitions). Now, applying proposition 4.13 to the simplicial complex $K(\Delta_\sigma)$ we get a cellular decomposition of a $(\text{codim } \sigma)$ -dimensional ball B_{Δ_σ} which realizes the simplicial complex $K(\Delta_\sigma)$. We have two geometric realizations of isomorphic simplicial complexes, so they must be isomorphic, and we conclude that $e_\sigma^\vee$ is homeomorphic to a $(\text{codim } \sigma)$ -dimensional ball.

It is clear that the cells $e_\sigma^\vee$ cover all the ball B , and that they are attached properly, that is the boundary of every cell is contained in a lower dimensional skeleton. ●

Remark 4.16. The cellular decomposition $B_\Delta^\vee$ is dual to the cellular decomposition of the sphere S_Δ together with an extra n -cell for the interior of the ball.

Proposition 4.17. There is a canonical isomorphism of chain complexes

$$\check{C}_*(\Delta, R) \simeq C_*^{\text{cell}}(B_\Delta^\vee, R).$$

Proof. There is a canonical isomorphism between the R -modules $\check{C}_k(\Delta, R)$ and $C_k^{\text{cell}}(B_\Delta^\vee, R)$, as both are generated by the cones in codimension k . Note that the unique n -cell in B_Δ corresponds to the zero cone in Δ .

It only remains to check that the differentials in $\check{C}_*(\Delta, R)$ coincide with the cellular ones. Note that the attaching maps $f_\tau: \partial e_\tau^\vee \rightarrow \text{Sk}_{\text{codim } \tau - 1} B_\Delta^\vee$ are homeomorphisms with the image. So, for any lower dimensional cell $e_\sigma^\vee$ on the boundary of $e_\tau^\vee$, the corresponding matrix element in the cellular differential is a sign, according to the relative orientation of the cells $e_\sigma^\vee$ and $e_\tau^\vee$. This is exactly the differential in $\check{C}_*(\Delta, R)$. ●

Corollary 4.18. *The augmented complex of sheaves*

$$\check{\mathcal{C}}_*(\Delta, R_X) \rightarrow R_X$$

is exact, that is, $\check{\mathcal{C}}_*(\Delta, R_X)$ is a resolution of the constant sheaf R_X .

Proof. From proposition 4.6 we know that the fiber of the complex $\check{\mathcal{C}}_*(\Delta, R_X) \rightarrow R_X$ at a point $x \in X(\Delta)$ is identified with $\check{C}(\Delta, R) \rightarrow R$. Now by proposition 4.17 we know that the complex $\check{C}(\Delta, R)$ is isomorphic to the cellular complex of the cellular decomposition of the ball $B_\Delta^\vee$, so its homology is

$$H_k \check{C}(\Delta, R) \simeq H_k C^{\text{cell}}(B_\Delta, R) = \begin{cases} R & \text{for } k = 0, \\ 0 & \text{for } k > 0. \end{cases}$$

Now, as the augmentation $a: \check{C}_*(\Delta, R) \rightarrow R$ is given by the sum of the elements on degree zero, we conclude that it is a quasi-isomorphism. $\bullet$

4.2 The spectral sequence

Let $X(\Delta)$ be a toric variety and $\mathcal{F}^*$ be a complex of sheaves on X . We describe a spectral sequence converging to the hypercohomology $\mathbb{H}^n(X(\Delta), \mathcal{F}^*)$.

Definition 4.19. A complex of sheaves $\mathcal{F}^*$ is said to have *homotopy invariant cohomology* if for every variety X the projection $p: X \times \mathbb{A}^1 \rightarrow X$ induces isomorphisms in hypercohomology

$$\mathbb{H}^n(X, \mathcal{F}^*) \xrightarrow{\sim} \mathbb{H}^n(X \times \mathbb{A}^1, \mathcal{F}^*).$$

Remark 4.20. The complex of sheaves $\mathcal{M}^*$ defining morphic cohomology has homotopy invariant cohomology by 2.26.

Theorem 4.21. *Let $X(\Delta)$ be a toric variety associated to a fan Δ and $\mathcal{F}^*$ a bounded above cochain complex of sheaves. There is a convergent spectral sequence*

$$E_1^{r,s} = \text{Ext}^r(\check{\mathcal{C}}_s(\Delta, \mathbb{Z}_X), \mathcal{F}^*) \implies \mathbb{H}^{r+s}(X(\Delta), \mathcal{F}^*). \quad (47)$$

Moreover, if $\mathcal{F}^*$ has homotopy invariant cohomology,

$$E_1^{r,s} = \bigoplus_{\sigma \in \Delta^{(s)}} \mathbb{H}^r(T_\sigma, \mathcal{F}^*), \quad (48)$$

and the differentials on the first page $d_1: E_1^{r,s} \rightarrow E_1^{r,s+1}$ are given by

$$d_1 = \sum_{\substack{\sigma \in \Delta^{(s)} \\ \tau \in \Delta^{(s+1)} \\ \tau \leq \sigma}} \epsilon(\tau, \sigma) r_{\tau, \sigma}^* \quad (49)$$

where

$$r_{\tau,\sigma}: T_\tau = \operatorname{Spec} \mathbb{C}[\tau^\perp] \rightarrow T_\sigma = \operatorname{Spec} \mathbb{C}[\sigma^\perp],$$

are the morphisms induced by the natural inclusion $\sigma^\perp \rightarrow \tau^\perp$.

Proof. Let $\mathcal{F}^* \rightarrow \mathcal{I}^*$ be a K-injective resolution of $\mathcal{F}^*$ (see remark 2.20). Let $\check{\mathcal{C}}_*(\Delta, \mathbb{Z}_X) \rightarrow \mathbb{Z}_X$ be the resolution of the constant sheaf $\mathbb{Z}_X$ from corollary 4.18. We build a double complex

$$C^{r,s} = \operatorname{Hom}(\check{\mathcal{C}}_s(\Delta, \mathbb{Z}_X), \mathcal{I}^r),$$

with the induced differentials (going in the increasing direction of r and s). The homology of this double complex in the r direction is $\operatorname{Ext}^r(\check{\mathcal{C}}_s(\Delta, \mathbb{Z}_X), \mathcal{F}^*)$, giving the spectral sequence

$$E_1^{r,s} = \operatorname{Ext}^r(\check{\mathcal{C}}_s(\Delta, \mathbb{Z}_X), \mathcal{F}^*) \implies \mathbb{H}^{r+s}(X(\Delta), \mathcal{F}^*).$$

As for the convergence. The complex of sheaves $\mathcal{I}^*$ is bounded above, and the schemes X_σ have finite cohomological dimension. Using the hypercohomology spectral sequence we conclude that $\mathbb{H}^k(X_\sigma, \mathcal{F}^*)$ is bounded above. In other words, the first page is bounded above in the r direction. It is clearly bounded (from both sides) in the s direction, and this is enough to establish the convergence.

If $\mathcal{F}^*$ is homotopy invariant, as the immersion $T_\sigma \rightarrow X_\sigma$ are algebraic homotopy equivalences we get the expression (48).

Finally, the differentials on the first page are induced by the s -differentials in the double complex $C^{r,s}$, which are given by the formula

$$d_1 = \sum_{\substack{\sigma \in \Delta^{(s)} \\ \tau \in \Delta^{(s+1)} \\ \tau \leq \sigma}} \epsilon(\tau, \sigma) i_{\tau,\sigma}^*$$

where $i_{\tau,\sigma}: X_\tau \rightarrow X_\sigma$ is the inclusion. The formula (49) follows from the equation

$$r_{\tau,\sigma}^* = j_\tau^* i_{\tau,\sigma}^* r_\sigma^*,$$

and the fact that j_τ^* and r_σ^* are mutually inverse isomorphisms giving the identification $\mathbb{H}^r(T_\sigma, \mathcal{F}^*) \simeq \mathbb{H}^r(X_\sigma, \mathcal{F}^*)$. ●

4.3 Some applications

We have a rather explicit description of the first page and differentials of the spectral sequence in 4.21. Together with the computation 3.18 of the morphic cohomology of a torus we can make it still more explicit.

Corollary 4.22. *Let $\mathcal{F}^* = \mathcal{M}^*$. Then the first page of the spectral sequence in 4.21 is*

$$E_1^{r,s} \simeq \bigoplus_{\sigma \in \Delta^{(s)}} \wedge^r(\sigma^\perp \cap M) \otimes LH(\text{pt}) \quad (50)$$

and the differentials $d_1^s: E_1^{r,s} \rightarrow E_1^{r,s+1}$ are given by

$$d_1^s \left(\sum_{\sigma \in \Delta^{(s)}} x_\sigma v_{1,\sigma} \wedge \cdots \wedge v_{r,\sigma} \right) = \sum_{\sigma \in \Delta^{(s)}} x_\sigma \sum_{\substack{\tau \in \Delta^{(s+1)} \\ \tau \leq \sigma}} \epsilon(\tau, \sigma) v_{1,\sigma} \wedge \cdots \wedge v_{r,\sigma}$$

Proof. This is a straight forward combination of theorem 4.21 and the computation 3.18. •

Finally, using an idea from [Jor98] which can be traced back to [Tot] that this spectral sequence degenerates rationally.

Theorem 4.23. *The spectral sequence in 4.22 degenerates when tensored with $\mathbb{Q}$.*

Proof. Let $LH(\text{pt})_{\mathbb{Q}} = LH(\text{pt}) \otimes_{\mathbb{Z}} \mathbb{Q}$. The toric variety $X(\Delta)$ admits an $\mathbb{N}$ -action. Let $m \in \mathbb{N}$, then $[m]: X(\Delta) \rightarrow X(\Delta)$ is the morphism which on the open sets $X_\sigma = \text{Spec } \mathbb{C}[\sigma^\vee \cap M]$ is defined through the ring homomorphism $[m]^*: \mathbb{C}[\sigma^\vee \cap M] \rightarrow \mathbb{C}[\sigma^\vee \cap M]$ given by $v \mapsto mv$ (see [Tot] for details).

The $\mathbb{N}$ -action on $X(\Delta)$ induces an $\mathbb{N}$ -action on the spectral sequence 50. As the rational morphic cohomology of a torus $T_L = \text{Spec } \mathbb{C}[L]$ is

$$LH^n(T_L)_{\mathbb{Q}} = \wedge^n L \otimes LH(\text{pt})_{\mathbb{Q}},$$

The $\mathbb{N}$ -action on the page $E_1^{r,s}$ is just multiplication by m^r . As the next pages $E_k^{r,s}$ of the spectral sequence are subquotients of $E_1^{*,*}$, the action on those pages is also given by m^r . On the other hand the differentials go $d_k: E_k^{r,s} \rightarrow E_k^{r+1-k,s+k}$ and the $\mathbb{N}$ -action commutes with them, so

$$m^r d_k(x) = d_k(m^r x) = d_k([m]x) = [m]d_k(x) = m^{r+1-k} d_k(x),$$

where $x \in E_k^{r,s}$. Rationally, this forces $d_k(x) = 0$ when $k \geq 2$. •

Now let $\varepsilon: \mathbf{Top} \rightarrow \mathbf{qProj}_{\mathbb{C}}$ be the morphism of sites, with the usual topology in $\mathbf{Top}$ and the Zariski topology on $\mathbf{qProj}_{\mathbb{C}}$. Let $\mathbf{R}\varepsilon_* \mathbb{Z}$ be the derived push-forward of the constant sheaf $\mathbb{Z}$ on $\mathbf{Top}$ to the Zariski site $\mathbf{qProj}_{\mathbb{C}}$. There is a natural map

$$\mathcal{M}^*(q) \rightarrow \mathbf{R}\varepsilon_* \mathbb{Z} \quad (51)$$

which, on smooth varieties, factors as

$$\mathcal{M}^*(q) \rightarrow \tau_{\leq q} \mathbf{R}\varepsilon_* \mathbb{Z}. \quad (52)$$

See [FHW04] for details.

There is the following conjecture, a morphic analogue of the Beilinson-Lichtenbaum conjecture in the motivic world.

Conjecture 4.24 (Suslin). *The comparison morphism (51) above is a quasi-isomorphism on smooth varieties.*

This conjecture is proved for the class of smooth linear varieties (which include smooth toric varieties) in [FHW04] theorem 7.14.

The spectral sequence 4.21 has the following corollary.

Corollary 4.25. *The Suslin conjecture holds for projective toric varieties (not necessarily smooth).*

Proof. First of all, we have to check that $\mathcal{M}^*(q)|_{X(\Delta)}$ is exact above degree q , in order to have the factorization (52). This is a local statment on $X(\Delta)$, so we can restrict to an open X_σ . Now the inclusion $T_\sigma \rightarrow X_\sigma$ is an algebraic homotopy equivalence, and they induce isomorphisms on hypercohomology

$$\mathbb{H}^n(X_\sigma, \mathcal{M}^*(q)|_X) \xrightarrow{\sim} \mathbb{H}^n(T_\sigma, \mathcal{M}^*(q)|_{T_\sigma}),$$

so the natural map $\mathcal{M}^*(q)|_X \rightarrow \mathbf{R}j_{\sigma*}j_\sigma^*\mathcal{M}^*(q)|_X$ is a quasi-isomorphism. As T_σ is smooth, its cohomology vanishes above q , and we have the desired factorization.

Now, $\tau_{\leq q}\mathbf{R}\mathcal{E}_*\mathbb{Z}$ has homotopy invariant cohomology. Then we can apply theorem 4.21 to it and get a spectral sequence converging to $\mathbb{H}^n(X(\Delta), \tau_{\leq q}\mathbf{R}\mathcal{E}_*\mathbb{Z})$. Moreover, the comparison map (52) gives a morphism of spectral sequences

$$\begin{array}{ccc} E_1^{r,s} = \bigoplus_{\sigma \in \Delta(s)} \mathbb{H}^r(T_\sigma, \mathcal{M}^*(q)) & \Longrightarrow & \mathbb{H}^{r+s}(X(\Delta), \mathcal{M}^*(q)) \\ \downarrow & & \downarrow \\ E_1'^{r,s} = \bigoplus_{\sigma \in \Delta(s)} \mathbb{H}^r(T_\sigma, \tau_{\leq q}\mathbf{R}\mathcal{E}_*\mathbb{Z}) & \Longrightarrow & \mathbb{H}^{r+s}(X(\Delta), \tau_{\leq q}\mathbf{R}\mathcal{E}_*\mathbb{Z}) \end{array}$$

which is an isomorphism on the first page by the computation 3.18, so it gives an isomorphism on the abutement, as desired. ●

References

- [DT58] Albrecht Dold and René Thom. Quasifaserungen und unendliche symmetrische Produkte. *Ann. of Math. (2)*, 67:239–281, 1958.
- [FG93] Eric M. Friedlander and Ofer Gabber. Cycle spaces and intersection theory. In *Topological methods in modern mathematics (Stony Brook, NY, 1991)*, pages 325–370. Publish or Perish, Houston, TX, 1993.
- [FHW04] Eric M. Friedlander, Christian Haesemeyer, and Mark E. Walker. Techniques, computations, and conjectures for semi-topological K -theory. *Math. Ann.*, 330(4):759–807, 2004.

-
- [FL92] Eric M. Friedlander and H. Blaine Lawson, Jr. A theory of algebraic cocycles. *Ann. of Math. (2)*, 136(2):361–428, 1992.
- [FL97] Eric M. Friedlander and H. Blaine Lawson. Duality relating spaces of algebraic cocycles and cycles. *Topology*, 36(2):533–565, 1997.
- [FM94] Eric M. Friedlander and Barry Mazur. Filtrations on the homology of algebraic varieties. *Mem. Amer. Math. Soc.*, 110(529):x+110, 1994. With an appendix by Daniel Quillen.
- [Fri91] Eric M. Friedlander. Algebraic cycles, Chow varieties, and Lawson homology. *Compositio Math.*, 77(1):55–93, 1991.
- [Fri98] Eric M. Friedlander. Algebraic cocycles on normal, quasi-projective varieties. *Compositio Math.*, 110(2):127–162, 1998.
- [Fri00] Eric M. Friedlander. Bloch-Ogus properties for topological cycle theory. *Ann. Sci. École Norm. Sup. (4)*, 33(1):57–79, 2000.
- [Ful93] William Fulton. *Introduction to toric varieties*, volume 131 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1993. The William H. Roever Lectures in Geometry.
- [FW01] Eric M. Friedlander and Mark E. Walker. Function spaces and continuous algebraic pairings for varieties. *Compositio Math.*, 125(1):69–110, 2001.
- [GJ99] Paul G. Goerss and John F. Jardine. *Simplicial homotopy theory*, volume 174 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 1999.
- [Jor98] A. Jordan. *Homology and cohomology of toric varieties*. PhD thesis, Universität Konstanz, 1998. Available from: <http://www.inf.uni-konstanz.de/Preprints/papers/1998/preprint-057.ps>.
- [Jos01] Roy Joshua. Algebraic K -theory and higher Chow groups of linear varieties. *Math. Proc. Cambridge Philos. Soc.*, 130(1):37–60, 2001.
- [Kol96] János Kollár. *Rational curves on algebraic varieties*, volume 32 of *Ergebnisse der Mathematik*. Springer-Verlag, Berlin, 1996.
- [Law89] H. Blaine Lawson, Jr. Algebraic cycles and homotopy theory. *Ann. of Math. (2)*, 129(2):253–291, 1989.
- [LF92] Paulo Lima-Filho. Lawson homology for quasiprojective varieties. *Compositio Math.*, 84(1):1–23, 1992.
- [MVW06] Carlo Mazza, Vladimir Voevodsky, and Charles Weibel. *Lecture notes on motivic cohomology*, volume 2 of *Clay Mathematics Monographs*. American Mathematical Society, Providence, RI, 2006.

-
- [Spa88] N. Spaltenstein. Resolutions of unbounded complexes. *Compositio Math.*, 65(2):121–154, 1988.
- [Tot] B. Totaro. Chow groups, Chow cohomology and linear varieties. preprint. Available from: <http://www.dpmms.cam.ac.uk/~bt219/linear.dvi>.
- [Voe00] V. Voevodsky. Homotopy theory of simplicial sheaves in completely decomposable topologies. preprint, 2000. Available from: <http://www.math.uiuc.edu/K-theory/0443/>.
- [Voi08a] Mircea Voineagu. Private communication, 2008.
- [Voi08b] Mircea Voineagu. Semi-topological K -theory for certain projective varieties. *J. Pure Appl. Algebra*, 212(8):1960–1983, 2008.

Abdó Roig-Maranges
abdo.roig@upc.edu

Universitat Politècnica de Catalunya
Dept. de Matemàtica Aplicada 1
Av. Diagonal, 647
08028 Barcelona (Spain)